

# Counter-regulatory renin–angiotensin system in cardiovascular disease

Maria Paz Ocaranza<sup>1,6</sup>, Jaime A. Riquelme<sup>2,6</sup>, Lorena García<sup>2</sup>, Jorge E. Jalil<sup>1</sup>, Mario Chiong<sup>2</sup>, Robson A. S. Santos<sup>3</sup> and Sergio Lavandero<sup>1,2,4,5\*</sup>

**Abstract** | The renin–angiotensin system is an important component of the cardiovascular system. Mounting evidence suggests that the metabolic products of angiotensin I and II — initially thought to be biologically inactive — have key roles in cardiovascular physiology and pathophysiology. This non-canonical axis of the renin–angiotensin system consists of angiotensin 1–7, angiotensin 1–9, angiotensin-converting enzyme 2, the type 2 angiotensin II receptor (AT<sub>2</sub>R), the proto-oncogene Mas receptor and the Mas-related G protein-coupled receptor member D. Each of these components has been shown to counteract the effects of the classical renin–angiotensin system. This counter-regulatory renin–angiotensin system has a central role in the pathogenesis and development of various cardiovascular diseases and, therefore, represents a potential therapeutic target. In this Review, we provide the latest insights into the complexity and interplay of the components of the non-canonical renin–angiotensin system, and discuss the function and therapeutic potential of targeting this system to treat cardiovascular disease.

The renin–angiotensin system (RAS) has a critical role in cardiovascular physiology through its effects in regulating blood pressure and electrolyte balance<sup>1</sup>. However, under pathophysiological conditions, the effects of the RAS can intensify to trigger inflammation and structural remodelling, thus promoting cardiac and vascular damage<sup>2,3</sup>. Researchers have studied the RAS for more than a century, not only to understand its role in normal physiological function but also to develop effective therapies to treat its dysregulation<sup>1,2</sup>. These systematic research efforts have led to the discovery of a non-canonical RAS, which has challenged the hypothesis that the RAS can only exert deleterious effects on the cardiovascular and renal systems. In the classical system, renin cleaves angiotensinogen to form angiotensin I, which is subsequently converted to angiotensin II by angiotensin-converting enzyme (ACE) (FIG. 1). Conversely, ACE2 can cleave angiotensin II to produce angiotensin 1–7, and can cleave angiotensin I to generate angiotensin 1–9<sup>1,3</sup>. Increasing evidence supports the concept that these systems work to produce opposite effects, suggesting a counter-balancing role for the two axes in cardiovascular physiology and disease. A timeline of key historical findings associated with the study and discovery of the counter-regulatory RAS is shown in BOX 1. In light of the emergence of multiple studies evaluating the effects and signalling pathways elicited by the counter-regulatory RAS in the past decade, we sought to provide an update on the current understanding of the complex regulation of the non-canonical RAS. In this Review, we discuss the

cardioprotective effects of the non-canonical RAS and provide a critical analysis of the current challenges that must be overcome to translate its therapeutic effects into the clinical context.

## Components of the counter-regulatory RAS Ligands

The counter-regulatory RAS is made up of various peptides, receptors and enzymes (FIG. 1). Whereas the effects of angiotensin 1–7 and angiotensin 1–9 on the cardiovascular system have been explored previously<sup>3</sup>, the potential roles of other counter-regulatory RAS components remain poorly understood. These non-canonical RAS components include alamandine, angiotensin 1–12 and angiotensin 1–5, as well as angiotensin 2–8 and angiotensin 3–8, which are also known as angiotensin III and IV, respectively<sup>3</sup>. FIGURE 2 shows the molecular structures of these peptides.

In the past 10 years, new evidence has emerged about the signalling pathways triggered by the counter-regulatory RAS, revealing their role as potential therapeutic targets for cardiovascular disease (CVD). Angiotensin 1–7 can act as a  $\beta$ -arrestin-biased agonist of the type 1 angiotensin II receptor (AT<sub>1</sub>R) without activating the G<sub>q</sub> subunit. This mechanism might contribute to the anti-hypertrophic properties of angiotensin 1–7, given that neither activation of the AT<sub>1</sub>R nor the proto-oncogene Mas receptor antagonists prevented the beneficial effects of this peptide<sup>4</sup>. Alamandine activates the AMP-activated protein kinase (AMPK)–nitric oxide (NO)

<sup>1</sup>Advanced Center for Chronic Diseases (ACCDiS), Division de Enfermedades Cardiovasculares, Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile.

<sup>2</sup>Advanced Center for Chronic Diseases (ACCDiS), Facultad de Ciencias Químicas y Farmacéuticas, Universidad de Chile, Santiago, Chile.

<sup>3</sup>Department of Physiology and Biophysics, Institute of Biological Sciences, Federal University of Minas Gerais, Minas Gerais, Brazil.

<sup>4</sup>Department of Internal Medicine (Cardiology Division), University of Texas Southwestern Medical Center, Dallas, TX, USA.

<sup>5</sup>Advanced Center for Chronic Diseases (ACCDiS), Facultad de Medicina, Universidad de Chile, Santiago, Chile.

<sup>6</sup>These authors contributed equally: Maria Paz Ocaranza, Jaime A. Riquelme.

\*e-mail: [slavander@uchile.cl](mailto:slavander@uchile.cl)

<https://doi.org/10.1038/s41569-019-0244-8>

### Key points

- Chronic activation of the renin–angiotensin system (RAS) promotes cardiovascular damage, an effect that is antagonized by components of the counter-regulatory RAS.
- Components of the counter-regulatory RAS, including angiotensin 1–7, angiotensin 1–9, alamandine and their receptors have been found to be protective in multiple cardiovascular diseases, such as hypertension and heart failure.
- Numerous preclinical studies have demonstrated the beneficial effects of the counter-regulatory RAS, but clinical trials confirming these observations are still scarce.
- The challenges in quantitating angiotensin 1–7, angiotensin 1–9 and alamandine associated with their short plasma half-life and similarity in their molecular structures must be overcome before these peptides can be evaluated in the clinical setting.

pathway via the Mas-related G protein-coupled receptor member D (MRGD), which prevents angiotensin II-induced hypertrophy<sup>5</sup>. By contrast, angiotensin 1–9 stimulates the AT<sub>2</sub>R–AKT signalling pathway to protect the myocardium against reperfusion-induced cell death<sup>6</sup>. Moreover, angiotensin 1–12 has been shown to regulate intracellular calcium transients and left ventricular contractile function in both normal rats and rats with heart failure (HF) via a chymase-dependent and cyclic AMP-dependent mechanism<sup>7</sup>. Additionally, angiotensin 1–5 has been found to induce atrial natriuretic peptide (ANP) secretion from isolated perfused rat atria by binding to the Mas receptor and activating the phosphatidylinositol 3-kinase–AKT–endothelial NO synthase pathway<sup>8</sup>. Lastly, angiotensinogen is the precursor for the entire RAS family of peptides, but, to date, no studies have shown that angiotensinogen can elicit direct biological effects on the heart. Nonetheless, the aryl hydrocarbon receptor nuclear translocator-like protein 1 has been shown to modulate blood pressure through a mechanism involving transcriptional regulation of angiotensinogen in a circadian manner in perivascular adipose tissue, which in turn increases local angiotensin II production<sup>9</sup>. These novel findings shed light on the complex regulation of the classical RAS and suggest a similar complexity for its counter-regulatory system. In this context, circadian expression of local angiotensinogen might affect organ-specific activity of peptides with known cardiovascular effects, such as angiotensin 1–7 or angiotensin 1–9, and requires further investigation.

### Receptors


In the non-canonical RAS, angiotensin 1–7 and angiotensin 1–9 bind to the Mas receptor and AT<sub>2</sub>R, respectively, whereas alamandine acts through the MRGD<sup>3</sup> (FIG. 3). Angiotensin 1–7 can also bind to the MRGD, but the functional relevance of this association remains unclear<sup>10</sup>. Emerging evidence reveals a more complex interaction between components of the classical and the counter-regulatory RAS than initially thought, given that angiotensin 1–7 has also been shown to bind to AT<sub>2</sub>R<sup>11</sup>. Moreover, AT<sub>1</sub>R can form heterodimers with the Mas receptor, which inhibits the activity of AT<sub>1</sub>R<sup>12</sup>. Using radiolabelling and dynamic mass redistribution experiments in cells overexpressing the Mas receptor, Gaidarov and colleagues found that although angiotensin 1–7 can antagonize angiotensin II signalling, angiotensin 1–7

does not bind directly to the Mas receptor<sup>13</sup>. These data conflict with an earlier study that demonstrated binding of fluorescent or <sup>125</sup>I-labelled angiotensin 1–7 to the Mas receptor<sup>14</sup>. Gaidarov and colleagues noted that in the absence of the Mas receptor, angiotensin 1–7 has no effect on angiotensin II signalling<sup>13</sup>. However, the investigators also reiterated that rigorously controlled experiments demonstrating interactions between angiotensin 1–7 and the Mas receptor are very scarce. Moreover, given that their findings suggest that angiotensin 1–7 does not bind to the Mas receptor, the researchers hypothesized that any cardioprotective effects of angiotensin 1–7 might be attributable to antagonism of angiotensin II signalling<sup>13</sup>. Nevertheless, additional studies are required to confirm whether angiotensin 1–7 is an endogenous agonist of the Mas receptor.

Meems and colleagues designed and synthesized NPA7, a peptide that can simultaneously activate the Mas receptor and the particulate guanylyl cyclase A receptor<sup>15</sup>. NPA7, generated by the fusion of angiotensin 1–7 with a 22-amino acid sequence of the B-type natriuretic peptide (BNP), reduced blood pressure, cardiac unloading and systemic vascular resistance, and exerted a more potent natriuretic and diuretic effect than separate administration of BNP and angiotensin 1–7. These observations raise the possibility that fusion of other counter-regulatory RAS ligands that can target more than one receptor might also induce a synergistic effect to mediate potent cardioprotective benefits. AT<sub>2</sub>R can form functional heterodimers with Mas receptors, highlighting the possibility of developing drugs that can selectively target monomers or oligomers to upregulate or downregulate specific cell signalling cascades in the cardiovascular system<sup>16</sup>. In addition, the crystal structures of human AT<sub>2</sub>R bound to a selective ligand indicate that the ligand can induce an active conformation of the receptor, suggesting that AT<sub>2</sub>R does not bind to G proteins or  $\beta$ -arrestins<sup>17</sup>. Tetzner and colleagues showed that angiotensin 1–7 can bind to the MRGD and that the AT<sub>2</sub>R antagonist PD123319 can block both the Mas receptor and MRGD<sup>10</sup>. This latter finding is particularly important, given the large number of studies utilizing PD123319 to assess the effects of AT<sub>2</sub>R activation. FIGURE 3 provides an overview of the signalling pathways triggered by the counter-regulatory RAS ligands upon binding to their receptors.

### Regulatory enzymes

ACE inhibitors are a first-line pharmacological therapy in the management of hypertension. Other proteases such as ACE2 and neprilysin (also known as neutral endopeptidase) have been identified as novel therapeutic targets, given that these enzymes can also reduce blood pressure. ACE2 might reduce blood pressure levels by generating angiotensin 1–7 from angiotensin II, whereas inhibition of neprilysin increases ANP levels<sup>18</sup>. In addition, the endogenous metabolic regulator fibroblast growth factor 21 (FGF21) can promote ACE2 generation in adipocytes and renal cells, thereby promoting the cleavage of angiotensin II to form angiotensin 1–7, suggesting that FGF21 can reduce angiotensin II-induced hypertension<sup>19</sup>.




**Fig. 1 | Classical and counter-regulatory renin–angiotensin pathways.** In the classical system, renin cleaves angiotensinogen to produce angiotensin I. This peptide can be processed by angiotensin-converting enzyme (ACE) to form angiotensin II, which in turn can bind to the type 1 angiotensin II receptor (AT<sub>1</sub>R) and AT<sub>2</sub>R<sup>3</sup>. AT<sub>1</sub>R activation increases aldosterone<sup>165</sup> and anti-diuretic hormone (ADH)<sup>166</sup> production, sympathetic nervous system (SNS) tone<sup>167</sup>, blood pressure<sup>168</sup>, vasoconstriction<sup>169</sup>, cardiac hypertrophy<sup>170</sup>, fibrosis<sup>171</sup>, inflammation<sup>172</sup>, vascular smooth muscle cell (VSMC) dedifferentiation<sup>173</sup> and reactive oxygen species (ROS) production<sup>36</sup>, while decreasing parasympathetic nervous system (PSNS) tone<sup>174</sup>, baroreflex sensitivity<sup>175</sup>, nitric oxide (NO) production<sup>176</sup> and natriuresis<sup>177</sup>. Angiotensin II can be further processed by aminopeptidase A (APA) to form angiotensin III, which also acts through AT<sub>1</sub>R. Angiotensin III can be cleaved by alanyl aminopeptidase N (APN) to generate angiotensin IV, which binds to AT<sub>4</sub>R, producing cardioprotective effects<sup>178</sup>, increasing natriuresis<sup>179</sup> and NO production<sup>180</sup>, as well as reducing vasoconstriction<sup>181</sup>, inflammation<sup>178</sup> and VSMC dedifferentiation<sup>182</sup>. Angiotensin I can also be cleaved by ACE2 and neprilysin (NEP) to produce angiotensin 1–9 and angiotensin 1–7, respectively<sup>3</sup>. Angiotensin 1–9 can activate AT<sub>2</sub>R to trigger natriuresis<sup>183</sup> and NO production<sup>73</sup>, thus mediating vasodilatory effects<sup>73</sup> and reducing blood pressure<sup>73</sup>. In addition, this peptide is cardioprotective<sup>6</sup> and can attenuate inflammation<sup>73</sup>, cardiac hypertrophy<sup>135</sup> and fibrosis<sup>73</sup>. Angiotensin 1–7 binds to the proto-oncogene Mas receptor (MasR) and reduces both blood pressure<sup>184</sup> and noradrenaline release in hypertensive rodents<sup>185</sup>. Conversely, activation of MasR increases NO generation<sup>186</sup>, natriuresis<sup>187</sup>, vasodilatation<sup>186</sup>, PSNS tone and baroreflex sensitivity<sup>188,189</sup>. Angiotensin 1–7 can also be formed from angiotensin II cleavage by ACE2 and be further metabolized to alamandine. Alternatively, angiotensin II can be processed by aspartate decarboxylase (AD) to produce angiotensin A, which can be converted to alamandine by ACE2. Upon binding to the Mas-related G protein-coupled receptor member D (MRGD), alamandine can promote the same effects reported for angiotensin 1–7<sup>5,67,190</sup>, with the exception of natriuresis. RAS, renin–angiotensin system.

Box 1 | Timeline of the discoveries related to the counter-regulatory RAS

The timeline in the figure shows a historical perspective of the most important findings associated with the counter-regulatory renin–angiotensin system (RAS). The proto-oncogene Mas receptor (MasR) was initially described as an oncogene and detected through its tumorigenicity in nude mice<sup>147</sup>. Angiotensin 1–9 and angiotensin 1–7 were first identified from hydrolytic cleavage of angiotensin I, and angiotensin I or angiotensin II, respectively<sup>148,149</sup>. In 1989, angiotensin 1–7 was found to have anti-hypertensive effects in rats upon unilateral injection into the medial “nucleus of the solitary tract” and into the dorsal motor nucleus of the vagus<sup>150</sup>. The earliest patents related to the components of the counter-regulatory RAS described the use of the MasR in an assay system for detecting angiotensin-blocking activity, a cDNA encoding the type 2 angiotensin II receptor (AT<sub>2</sub>R) in mice and rats, a nucleic acid encoding angiotensin 1–7, a cDNA encoding angiotensin-converting enzyme 2

(ACE2) and the use of angiotensin 1–9 to prevent, reverse, inhibit or reduce cardiovascular, pulmonary, cerebral or renal remodelling. ACE2 was simultaneously discovered by two independent research groups in 2000<sup>151,152</sup>. Angiotensin 1–7 was subsequently described as a cardioprotective peptide<sup>153</sup> with anti-inflammatory actions<sup>110</sup> and found to be activated through the MasR<sup>14</sup>. The first clinical trial of angiotensin 1–7 assessed the effect of this peptide on the reduction of blood flow in solid tumours<sup>154</sup>, whereas the first trial of ACE2 evaluated the safety and tolerability of a recombinant form of ACE2<sup>121</sup>. The anti-hypertrophic<sup>135</sup>, anti-hypertensive<sup>73</sup>, anti-inflammatory<sup>120</sup> and cardioprotective<sup>6</sup> properties of angiotensin 1–9 have been described. Alamandine was discovered in 2013 as an anti-hypertensive agent<sup>67</sup>, and the cardioprotective properties of this compound have since been described<sup>155,156</sup>.



I–R, ischaemia–reperfusion.

**Intercellular communication**

The classical RAS can act at both local and systemic levels, but how these signals are coordinated is poorly understood. Exosomes, which are extracellular vesicles of 50–100 nm in size, can transport and transmit molecules such as proteins and microRNAs from one cell to another, and can also transport components of the classical RAS<sup>20</sup>. Previously assumed to be scattered cellular waste, exosomes are attracting much research interest since the discovery of their role in intercellular communication<sup>21</sup>. Considering that these extracellular vesicles can communicate signals from afar and that the counter-regulatory RAS can exert its effects on multiple cell types, these vesicles might have a role in orchestrating the effects of the counter-regulatory RAS. In this context, Pironti and colleagues observed that exosomes induced by cardiac pressure overload in mice contain functional AT<sub>1</sub>R, which might influence AT<sub>1</sub>R-mediated regulation of vascular tone<sup>22</sup>. Moreover, exosomes seem to have a role in the local RAS. Angiotensin II triggers exosome production in rat cardiac fibroblasts in vitro, and these exosomes in turn promote angiotensin II production and AT<sub>1</sub>R expression in rat cardiomyocytes in vitro, suggesting a positive feedback mechanism that might contribute to the exacerbation of cardiac hypertrophy elicited by angiotensin II<sup>23</sup>. However, this evidence only supports a role for exosomes in orchestrating the effects of the canonical RAS. Whether extracellular

vesicles contribute to the cardioprotective properties of the counter-regulatory RAS remains to be determined.

**Counter-regulatory RAS in CVD  
Pulmonary arterial hypertension**

**The ACE2–angiotensin 1–7–Mas receptor axis.** ACE2, first described as a receptor for severe acute respiratory syndrome coronavirus, is characterized by its marked homology with ACE<sup>24</sup>. The therapeutic potential of ACE2 agonists for pulmonary arterial hypertension (PAH) has been explored in a number of studies. In rats with monocrotaline-induced PAH, *Ace2* gene therapy prevented PAH-mediated hypertrophy and functional impairment of the right ventricle<sup>25</sup>. Moreover, synthetic activators of ACE2 (XNT<sup>26</sup> and resorcinolnaphthalen<sup>27</sup>) improve pulmonary artery endothelial function by inducing phosphorylation of endothelial NO synthase at Ser1177 and dephosphorylation at Thr495<sup>27</sup>, which consequently increases the bioavailability of NO. A meta-analysis to assess the efficacy of 522 interventions for PAH revealed that these ACE2 synthetic activators were among the most potent agents<sup>28</sup>. Although these findings strongly support the therapeutic potential of ACE2 activators, translation of these agents into a clinical setting remains challenging because ACE2 is a membrane-bound enzyme. ACE2 can be cleaved and its soluble and catalytically active form can be secreted<sup>29,30</sup>. Given that increasing the circulating levels




Fig. 2 | **Molecular structures of peptides of the counter-regulatory RAS.** The separation of these peptides from a biological sample is difficult, given the similarity of their molecular structures. Angiotensin 1-7 is only two amino acids shorter than angiotensin 1-9, and angiotensin 1-7 and alamandine only differ in their N-terminal amino acid. RAS, renin-angiotensin system.


of ACE2 might have a therapeutic effect, a recombinant human ACE2 (rhACE2) has been developed and tested in animal models. Administration of rhACE2 improved right ventricular function in mice subjected to pressure overload<sup>31</sup> and attenuated vascular remodelling in mice with bleomycin-induced pulmonary hypertension<sup>32</sup>. A pilot study evaluated the effects of increasing the enzymatic activity of ACE2 through intravenous infusion of 0.2 mg/kg or 0.4 mg/kg of rhACE2 in patients with PAH<sup>33</sup>. The drug was well tolerated and had beneficial effects on pulmonary vascular resistance and cardiac output, in addition to reducing inflammatory markers and increasing superoxide dismutase 2 levels in plasma. Nonetheless, this proof-of-concept study included only five patients. In a separate study, rhACE2 administration was also shown to be well tolerated in 44 patients with acute respiratory distress syndrome<sup>34</sup>. The safety profile of rhACE2 needs to be further assessed in clinical studies.

Angiotensin 1-7 and other Mas receptor activators might also have a protective role against the development of PAH<sup>35</sup>. Notably, however, angiotensin 1-7 is not considered a good therapeutic candidate owing to its pharmacokinetic limitations. Angiotensin 1-7 is rapidly cleaved by peptidases and thus has a very short half-life of ~10 s (REF.<sup>36</sup>). However, cell signalling mechanisms and

effects mediated by biological peptides are thought to persist despite their short half-life<sup>37</sup>. Furthermore, studies in animal models have shown that administration of angiotensin 1-7 included in cyclodextrin complexes has neuroprotective effects and improves muscle damage induced by eccentric cardiac overload<sup>38-44</sup>. A stable, cyclic analogue of angiotensin 1-7 moderately reduced right ventricular systolic pressure in a rat model of monocrotaline-induced PAH, but no significant changes were observed in the medial wall thickness of pulmonary arterioles<sup>45</sup>. To optimize the protective potential of this angiotensin 1-7 analogue for the treatment of PAH, the compound can potentially be combined with a neprilysin inhibitor or an ACE2 activator<sup>46</sup>; whether this approach is effective in maintaining high levels of angiotensin 1-7 requires further investigation.

**AT<sub>2</sub>R stimulation.** AT<sub>2</sub>R activation can attenuate right ventricular and pulmonary remodelling<sup>47</sup>. AT<sub>2</sub>R stimulation protected mice from severe lung injury induced by sepsis or acid aspiration<sup>48</sup>, whereas AT<sub>2</sub>R deficiency exacerbated HF in mice subjected to acute myocardial infarction<sup>49</sup>. Furthermore, activation of AT<sub>2</sub>R (using the agonist dKc-angiotensin 1-7) in a rat model of chronic lung disease protected the heart and lungs from





**Fig. 3 | Signal transduction mechanisms of the counter-regulatory RAS.** Signalling through the type 2 angiotensin II receptor (AT<sub>2</sub>R) can directly inhibit AT<sub>1</sub>R activation and thus antagonize the effects of angiotensin II<sup>191</sup>. Stimulation of AT<sub>2</sub>R can also inhibit extracellular signal-regulated kinase 1 (ERK1) and ERK2 by activating Src homology region 2 domain-containing phosphatase 1 (SHP1)<sup>192</sup> and mitogen-activated protein kinase-phosphatase 1 (MKP1)<sup>193</sup>, which can result in attenuation of cardiac hypertrophy. AT<sub>2</sub>R can also activate the transcription factor promyelocytic zinc finger protein (PLZF), thereby inducing the expression of ribosomal protein S6 kinase β1 (p70S6K) and p85α expression and, in turn, eliciting protein synthesis<sup>194</sup>. In addition, AT<sub>2</sub>R might trigger vasodilatation by activating the phosphatidylinositol-3-kinase (PI3K)–AKT–endothelial nitric oxide synthase (eNOS)–nitric oxide (NO)–cGMP pathway either via angiotensin 1–9-mediated activation<sup>194–196</sup> or by heterodimerization with bradykinin B<sub>2</sub> receptor (B<sub>2</sub>R)<sup>197</sup>. Phosphorylation of AKT by activation of AT<sub>2</sub>R through angiotensin 1–9 binding has also been found to confer cardioprotection<sup>6</sup>. Angiotensin 1–7 might induce the NO–soluble guanylyl cyclase pathway, thereby triggering vasodilatation via proto-oncogene Mas receptor (MasR) activation. Activation of this receptor can also reduce cardiac fibrosis by stimulating SHP1<sup>198</sup> and dual-specificity phosphatase (DUSP)<sup>199</sup>, consequently inhibiting p38 mitogen-activated protein kinase (MAPK) and ERK1 and ERK2<sup>200</sup>. The K<sub>Ca</sub>3.1 channel<sup>201</sup> and mothers against decapentaplegic homologue 2 (SMAD2) and SMAD3<sup>202</sup> are downstream targets of ERK1 and ERK2, and are downregulated upon MasR activation. Additionally, angiotensin 1–7 exerts an anti-hypertrophic effect by inhibiting nuclear factor of activated T cells (NFAT) through a MasR–PI3K–AKT–NO–cGMP-dependent pathway<sup>203</sup>. This anti-hypertrophic effect also depends on atrial natriuretic peptide (ANP) secretion during atrial pacing and is associated with activation of the Na<sup>+</sup>/H<sup>+</sup> exchanger (NHE1) and calcium/calmodulin-dependent protein kinase II (CaMKII) via the PI3K–AKT pathway<sup>200</sup>. Cardiac hypertrophy can also be reduced by activation of the Mas-related G protein-coupled receptor member D (MRGD) by alamandine via adenylate cyclase (AC)–cAMP–protein kinase A (PKA) signalling<sup>10</sup>.

damage by diminishing the inflammatory response and attenuating right ventricular hypertrophy, as well as reducing vascular wall thickness and alveolar septum thickness<sup>50</sup>. The AT<sub>2</sub>R agonist compound 21 (C21) has also been shown to inhibit cardiopulmonary fibrosis and right ventricular remodelling in a rat model of monocrotaline-induced PAH<sup>49</sup>. To date, only one pre-clinical study has assessed the effect of angiotensin 1–9 on PAH<sup>48</sup>. Adult rats with PAH treated with angiotensin 1–9 showed reduced right ventricular weight and systolic pressure, as well as diminished lung fibrosis, pulmonary arteriole thickness and endothelial damage compared with untreated controls. These effects were dependent on activation of the AT<sub>2</sub>R but not the Mas

receptor. Treatment with angiotensin 1–9 also reduced plasma levels of the pro-inflammatory markers tumour necrosis factor (TNF), CC-chemokine ligand 2 (CCL2; also known as MCP1), IL-1β and IL-6<sup>48</sup>.

**Systemic hypertension and remodelling**

**The ACE2–angiotensin 1–7 axis.** Numerous preclinical studies have shown that stimulating ACE2 with synthetic activators (such as XNT<sup>51</sup> and diminazene aceturate (DIZE))<sup>52</sup>, Mas receptor agonists such as AVE0991<sup>53</sup>, CGEN-856S<sup>54</sup> and CGEN-857<sup>54</sup> and human recombinant ACE2<sup>55</sup> can reduce blood pressure and attenuate cardiovascular damage. However, others studies have not found an association between hypertension and

ACE2 activity. The synthetic ACE2-activator XNT reduced blood pressure in an angiotensin II-induced model of hypertension, but plasma concentrations of angiotensin II and angiotensin 1–7 remained unaltered<sup>56</sup>. Moreover, the antihypertensive effect of this drug was observed in ACE2-deficient mice, and neither XNT nor DIZE induced the enzymatic activity of ACE2 in rat or mouse kidneys<sup>56</sup>. These findings raise the question as to whether researchers should continue to focus on these drugs with unknown mechanisms of action. However, ACE2 remains an appealing therapeutic target for treating hypertension, especially in tissues in which expression of this enzyme is higher than in plasma<sup>56</sup>. The therapeutic potential of DIZE as an alternative treatment for hypertension and PAH has been shown in previous experimental studies<sup>52,57</sup>. Moreover, deoxycorticosterone acetate (DOCA)-salt hypertensive rats treated with the Mas receptor agonist AVE0991 had lower blood pressure levels than untreated controls<sup>58</sup>. The anti-hypertrophic effects of AVE0991 are, in part, mediated by inhibition of NADPH oxidase 2 and NADPH oxidase 4, as observed in hypertensive mice subjected to aortic banding<sup>59</sup>. At present, the effects of these ACE2 activators have only been evaluated in preclinical studies. A rigorous evaluation of how these agents exert their beneficial effects is needed before they can be tested in the clinical setting, in order to identify off-target and potentially toxic effects.

ACE2 activity has also been assessed in patients with high blood pressure. The level of ACE2-mediated angiotensin II-degrading activity in monocyte-derived macrophages *in vitro* has been found to be similar in cells from both healthy individuals and patients with hypertension<sup>60</sup>. Of note, ACE2 activity is significantly higher in monocyte-derived macrophages from patients with prehypertension than in those from patients with hypertension, suggesting a potential role for ACE2 as an early marker of hypertension. This finding might also indicate a physiological protective mechanism against hypertension, most probably through the rapid degradation of angiotensin II<sup>60</sup>. By contrast, no correlation has been found between hypertension and ACE2 activity in patients with ST-segment elevation myocardial infarction<sup>61</sup>.

Plasma ACE2 levels have been suggested to vary depending on sex<sup>62,63</sup>, although most of the research exploring the role of ACE2 in CVD has not considered sex-related differences in activity levels. During pregnancy, plasma levels of angiotensin II are significantly elevated, whereas angiotensin 1–7 levels are significantly diminished, which together might predispose pregnant women to hypertension-related complications<sup>64</sup>. Furthermore, levels of urinary angiotensin 1–7 in patients with hypertension have been reported to be inversely proportional to blood pressure levels, implying a crucial role for this peptide in the development of hypertension<sup>65</sup>. Finally, angiotensin 1–7 has also been shown to alleviate obesity-induced haemodynamic alterations<sup>66</sup>.

**Alamandine.** Alamandine is a heptapeptide formed by the catalytic action of ACE2 on angiotensin A or directly from angiotensin 1–7 in the heart. Oral administration of an inclusion compound of alamandine and  $\beta$ -hydroxypropyl cyclodextrin reduced blood pressure

in spontaneously hypertensive rats and diminished myocardial fibrosis in isoprenaline-treated rats<sup>67</sup>. This anti-hypertensive effect was shown to have two phases. Initially, mean arterial pressure and left ventricular systolic pressure increased briefly in an AT<sub>1</sub>R-dependent manner, followed by a reduction in these parameters, which persisted throughout the rest of the infusion period. This anti-hypertensive effect was reversed by PD123319, an AT<sub>2</sub>R antagonist<sup>68</sup>. Additionally, alamandine treatment mitigated vascular remodelling in mice subjected to transverse aortic constriction<sup>69</sup>. Additional studies are required to further our understanding of the complex regulation of alamandine, the cell signalling cascades it triggers, and its therapeutic implications for hypertension and other CVDs. The normal range of alamandine levels in both healthy individuals and patients with hypertension should be established to provide a better understanding of the effect of RAS inhibition on alamandine plasma concentrations in this clinical context.

**AT<sub>2</sub>R agonists.** The vasodilatory effects of AT<sub>2</sub>R activation have been demonstrated in mice lacking<sup>70,71</sup> or overexpressing this receptor<sup>72</sup>. Mice lacking the AT<sub>2</sub>R showed an increased response to angiotensin II and significantly elevated blood pressure levels<sup>70,71</sup>, whereas transgenic overexpression of the AT<sub>2</sub>R in vascular smooth muscle cells of mice reduced angiotensin II-induced vasoconstriction<sup>72</sup>. The anti-hypertensive effects of the AT<sub>2</sub>R-selective agonists CGP42112A and angiotensin 1–9 have also been evaluated<sup>73,74</sup>. CGP42112A-treated obese rats had reduced blood pressure levels compared with untreated rats, which was associated with an increase in urinary sodium excretion<sup>74</sup>. This agonist also decreased blood pressure levels in spontaneously hypertensive rats<sup>75</sup> and prevented endothelial cell migration mediated by vascular endothelial growth factor signalling<sup>76</sup>.

The specific Rho kinase inhibitor fasudil significantly increased plasma levels of angiotensin 1–9 in both normotensive and hypertensive rats<sup>77</sup>. In addition, fasudil reduced blood pressure levels and aortic Rho kinase and ACE activity, whereas mRNA and protein levels of ACE2 were increased in plasma and the aortic wall<sup>77</sup>. Interestingly, another study showed an increase in ACE and angiotensin II levels in patients at high risk of acute pulmonary embolism compared with healthy volunteers<sup>78</sup>. Moreover, in a rat model of acute pulmonary embolism, RhoA–ROCK signalling mediated an imbalance in RAS vasoconstrictors, which was reversed with ROCK inhibitors or an ACE2 activator<sup>78</sup>. These findings further highlight the protective effects that ROCK inhibition can exert in the setting of hypertension, atherosclerosis and pathological cardiovascular remodelling.

In a study by Ocaranza and colleagues, administration of angiotensin 1–9 reduced blood pressure levels in hypertensive rats and attenuated myocardial damage by inhibiting the development of ventricular hypertrophy and fibrosis; importantly, these effects were mediated through AT<sub>2</sub>R but not Mas receptor signalling<sup>73</sup>. However, in a separate study, gene delivery of angiotensin 1–9 with an adeno-associated virus (AAV) in mice subjected to coronary artery ligation completely restored

systolic blood pressure levels and cardiac output compared with sham-treated mice, but histological analysis revealed only mild effects on cardiac hypertrophy and fibrosis<sup>79</sup>. Notably, Ocaranza and colleagues only evaluated angiotensin 1–9 administration for 2 weeks<sup>73</sup>, compared with the latter study that examined the effects of this peptide for 8 weeks<sup>79</sup>. The conflicting findings between these two studies suggest that the attenuation of myocardial damage might be transient and not sustained in the long term. However, the latter study did not measure plasma levels of angiotensin 1–9. AAV-mediated gene delivery of angiotensin 1–9 might not have produced a therapeutic concentration of the peptide in the blood that would be sufficient to protect the heart from adverse structural remodelling. A study that tested the anti-hypertensive actions of angiotensin 1–9 in stroke-prone spontaneously hypertensive rats also found no evidence of a protective effect<sup>80</sup>, but this study used a dose of angiotensin 1–9 that was six times lower than that used by Ocaranza and colleagues<sup>73</sup>. Additional studies are warranted to explore the anti-hypertensive and anti-remodelling effects of angiotensin 1–9 administration and the implications of the plasma levels of this peptide on cardioprotection. Although the efficacy of angiotensin 1–9 administration has not been explored in the clinical setting, in patients with acute respiratory distress syndrome, higher angiotensin 1–9 levels in plasma were associated with reduced mortality, whereas increased plasma angiotensin I levels were associated with increased mortality<sup>81</sup>.

### Heart failure

ACE2 is critical for heart function<sup>82</sup>, vasodilatation<sup>83</sup> and fluid balance<sup>84</sup>. *Ace2*<sup>-/-</sup> mutant mice have impaired contractility, increased expression of hypoxia markers and increased circulating levels of angiotensin II compared with control mice<sup>82</sup>. Furthermore, *Ace2*<sup>-/-</sup> mutant mice develop angiotensin II-mediated dilated cardiomyopathy that is characterized by an increase in markers of oxidative stress and inflammation, pathological hypertrophy and impaired left ventricular function<sup>85</sup>. Interestingly, plasma levels of the soluble form of ACE2 have been reported to be elevated in patients with HF and reduced ejection fraction, suggesting that sustained activation of the counter-regulatory RAS in HF might be a compensatory mechanism to attenuate cardiovascular dysfunction<sup>86</sup>. The mechanisms underlying HF with preserved ejection fraction (HFpEF) remain poorly defined, but the progression of this disease has been proposed to be linked to hypertension-induced cardiac remodelling<sup>87</sup>. Given the anti-hypertensive and anti-remodelling effects of the counter-regulatory RAS described thus far, this non-canonical signalling pathway might be a potential therapeutic target for the treatment of HFpEF. Angiotensin II infusion in wild-type mice resulted in increased blood pressure levels, myocardial hypertrophy, fibrosis and diastolic dysfunction; these effects were exacerbated in *Ace2*<sup>-/-</sup> mice<sup>88</sup>. Conversely, treatment of angiotensin II-infused wild-type mice with rhACE2 reduced angiotensin II-induced superoxide production and blunted the cardiac hypertrophic response, highlighting a possible protective role for this enzyme in HFpEF<sup>88</sup>.

Other components of the non-canonical RAS pathway are also involved in HF. Mice deficient in the alamandine receptor MRGD have left ventricular remodelling and severe dysfunction, and present with pronounced dilated cardiomyopathy<sup>89</sup>. Furthermore, infusion of the AT<sub>2</sub>R agonist C21 for 7 days in rats with HF induced by coronary artery ligation led to a reduction in noradrenaline excretion, as well as decreased renal sympathetic nerve activity<sup>90</sup>. Additionally, C21 administration increased baroreflex sensitivity, suggesting a protective role for this drug in the setting of HF.

Collectively, these findings support a role for various components of the counter-regulatory RAS in HF, both as potential biomarkers and therapeutic targets. Additional clinical studies are needed to determine the levels of ACE2, angiotensin 1–9 and angiotensin 1–7 in patients with HF.

### Myocardial infarction

The role of non-canonical RAS signalling in the development of myocardial infarction has been described. *ACE2* mRNA levels are elevated in the setting of myocardial infarction<sup>91</sup>, whereas loss of ACE2 can further exacerbate cardiac damage<sup>92</sup>. By the same token, *Ace2* overexpression has been shown to alleviate myocardial damage induced by ischaemia–reperfusion in rats<sup>93</sup>. Furthermore, administration of angiotensin 1–7 (added to the oligosaccharide hydroxypropyl  $\beta$ -cyclodextrin) in rats with myocardial infarction improved cardiac function and reduced infarct size by 50%<sup>42,43</sup>. Likewise, transgenic rats overexpressing a fusion protein that leads to a selective increase in angiotensin 1–7 levels were less susceptible to reperfusion-induced arrhythmias and isoproterenol-induced hypertrophy than wild-type rats<sup>94</sup>.

The cardioprotective role of AT<sub>2</sub>R in preventing post-ischaemic cardiac remodelling has been documented<sup>95,96</sup>. Mice lacking AT<sub>2</sub>R have aggravated myocardial infarction-induced HF and reduced survival compared with sham-treated mice<sup>97</sup>. Correspondingly, transgenic mice overexpressing AT<sub>2</sub>R showed improved left ventricular function after myocardial infarction<sup>98</sup>, and similar results were observed in rats with cardiac-specific overexpression of AT<sub>2</sub>R<sup>99</sup>. Administration of the AT<sub>2</sub>R agonist C21 to rats subjected to coronary artery ligation significantly improved recovery of left ventricular function and reduced cardiac remodelling after myocardial infarction<sup>100</sup>. Delivery of angiotensin 1–9 with an AAV vector into mice after the induction of myocardial infarction resulted in a reduction in sudden cardiac death and improved left ventricular function compared with control mice<sup>79</sup>. Importantly, angiotensin 1–9 had a positive inotropic effect, achieved by increasing calcium-transient amplitude and contractility through a protein kinase A-dependent mechanism<sup>79</sup>. Using an ex vivo approach with isolated rat hearts subjected to global ischaemia and reperfusion, Mendoza-Torres and colleagues showed that angiotensin 1–9 infusion can also reduce infarct size and apoptotic and necrotic cell death, and improve left ventricular function in an AT<sub>2</sub>R-dependent and AKT-dependent mechanism<sup>6</sup>. Together, these data suggest that angiotensin 1–7 and angiotensin 1–9



might be valuable pharmacological tools for the treatment of myocardial infarction, given their acute and long-term cardioprotective effects.

### Inflammation

Inflammatory processes are central to the development and progression of CVDs such as atherosclerosis, hypertension, myocardial infarction and HF<sup>101–105</sup>. A link between inflammation and RAS has previously been observed. T cells have an endogenous RAS that can regulate T cell function, NADPH oxidase activity and superoxide production<sup>106,107</sup>. Natural killer cells have also been shown to express renin, angiotensinogen, ACE and AT<sub>2</sub>R<sup>107</sup>. In line with these observations, the pro-inflammatory state is thought to upregulate RAS signalling in the setting of hypertension<sup>108</sup>. Interestingly, human monocytes also express ACE and ACE2 and can produce angiotensin 1–7 and angiotensin 1–9<sup>109</sup>. Taken together, these data suggest that the immune system might also be involved in regulating the non-canonical RAS.

Activation of the Mas receptor has been shown to promote anti-inflammatory effects<sup>110</sup>. Mice lacking this receptor have an exacerbated inflammatory reaction after treatment with lipopolysaccharides compared with wild-type mice<sup>111</sup>. Therefore, Mas receptor activation might be a valuable therapeutic target to counteract the pro-inflammatory processes that promote the development and progression of atherosclerosis<sup>112,113</sup>. Indeed, the Mas receptor agonist AVE0991 inhibits atherogenesis in *Apoe*<sup>-/-</sup> mice<sup>114</sup>. Moreover, long-term angiotensin 1–7 treatment confers both vasoprotection (by improving endothelial function) and atheroprotection (by reducing lesion progression) in *Apoe*<sup>-/-</sup> mice<sup>115</sup>. Consistent with these observations, angiotensin 1–7 can activate signalling pathways critical for the resolution of inflammatory processes involved in asthma<sup>116</sup>.

In addition to the Mas receptor, AT<sub>2</sub>R signalling has also been associated with the regulation of inflammation. The AT<sub>2</sub>R agonist C21 dose-dependently attenuates lipopolysaccharide-induced TNF and IL-6 production, but increased production of the anti-inflammatory cytokine IL-10<sup>117</sup>. Consistent with these observations, a separate study showed that administration of C21 in prehypertensive, obese Zucker rats reduced plasma levels of TNF and IL-6, whereas coadministration with the AT<sub>2</sub>R antagonist PD123319 decreased IL-10 levels in the kidneys<sup>118</sup>. Furthermore, in Wistar rats subjected to left coronary artery ligation, C21 treatment reduced the production of the pro-inflammatory cytokines IL-1 $\beta$ , IL-6 and IL-2 in an AT<sub>2</sub>R-dependent manner, improved systolic and diastolic ventricular function, and reduced scar size<sup>119</sup>. Angiotensin 1–9 administration has also been shown to reduce cardiac and renal inflammation in a DOCA–salt model of hypertension in rats, but this effect was independent of AT<sub>2</sub>R<sup>120</sup>.

### From bench to bedside

Research into the counter-regulatory RAS has resulted in the generation of a substantial amount of intellectual property related to its study and use. Currently, 184 patent applications associated with this system have been filed, most related to angiotensin 1–7 and its analogues,

AT<sub>2</sub>R, the Mas receptor, ACE2 and angiotensin 1–9. Only 76 patents are related to cardiovascular applications involving the control of arterial pressure, vascular remodelling, cardiac remodelling and HF. Furthermore, the robust evidence collated from large numbers of pre-clinical studies on the counter-regulatory RAS has also prompted the initiation of numerous clinical trials. At the time of this report, 15 clinical trials that involve interventions with counter-regulatory RAS molecules in CVDs were ongoing, including two studies designed to evaluate the safety of recombinant ACE2 and angiotensin 1–7 in treating thrombocytopenia<sup>121,122</sup>. A further nine trials aim to assess the use of ACE2 in the treatment of pulmonary hypertension<sup>123,124</sup> and the safety and use of angiotensin 1–7 in hypoxia, hypertension, HF and coronary artery bypass surgery<sup>125–131</sup>. Two trials investigating the use of angiotensin 1–7 to treat peripheral arterial disease and obesity-associated hypertension<sup>132,133</sup> are currently in the pre-recruitment phase.

### Challenges in interpretation

Despite the substantial amount of evidence suggesting a counter-regulatory role for the non-canonical RAS in protecting against the deleterious actions of a dysregulated classical RAS, the complexity of the relationship between the two systems remains to be fully elucidated. For example, ACE2 is elevated in patients with HF<sup>86</sup> or pre-hypertension<sup>60</sup>, but depressed in patients with PAH<sup>33</sup>. These discrepancies suggest that the components of the counter-regulatory RAS are upregulated or downregulated depending on the stage, severity or type of CVD. Moreover, these conflicting findings reinforce our lack of knowledge of the physiological and pathophysiological mechanisms involved in non-canonical RAS regulation. For instance, elevated levels of soluble ACE2 might represent a compensatory mechanism in response to HF but might also be the result of increased cleavage of membrane ACE2 by disintegrin and metalloproteinase domain-containing protein 17, which is known to be upregulated in HF<sup>86</sup>. In addition, RAS peptides can also be modulated by pharmacological treatment. In this regard, patients with chronic HF treated with ACE inhibitors have elevated plasma levels of angiotensin 1–7 and reduced plasma levels of angiotensin II, whereas patients with acute HF treated with angiotensin II receptor antagonists have decreased plasma levels of angiotensin 1–7 and increased plasma levels of angiotensin II<sup>134</sup>. Furthermore, the addition of rhACE2 to plasma samples from patients with HF induced the conversion of angiotensin I and angiotensin II into angiotensin 1–9 and angiotensin 1–7, respectively<sup>134</sup>.


### Limitations in application

In addition to the aforementioned challenges in interpreting the data on the non-canonical RAS, the measurement of angiotensin 1–7, angiotensin 1–9 or alamanidine in a clinical context poses many challenges. The separation of these peptides from a biological sample is difficult, given the similarity in their molecular structures. Angiotensin 1–7 is only two amino acids shorter than angiotensin 1–9<sup>135</sup>, whereas angiotensin 1–7 and alamanidine only differ in their N-terminal amino acid<sup>67</sup> (FIG. 2).

## Box 2 | Quantification of angiotensin peptides

Three critical issues should be considered to ensure that the method for quantifying angiotensin peptides is reliable. First, the sampling procedure must be efficient because blood or tissue samples need to undergo immediate peptidase inhibition to ensure stabilization of angiotensin metabolites<sup>157–162</sup> (see the figure, part a). The sampling duration should be kept as short as possible to avoid unexpected shifts in angiotensin metabolite patterns. Second, the structural similarity of all angiotensin metabolites (FIG. 2) necessitates an effective separation procedure, usually liquid chromatography<sup>157–162</sup>.

Once all peptides have been separated, angiotensin metabolites are quantified in the liquid chromatography eluate. Finally, the assay must be sensitive, given that angiotensin metabolites have been described at levels in the femtomolar range<sup>157–162</sup>. Immunological assays (radioimmunoassay (RIA) or enzyme-linked immunosorbent assays (ELISA))<sup>157–160</sup> and mass spectrometry<sup>161,162</sup> have been used in this setting. Both methods have lower limits of quantification for angiotensin metabolites (~1–2 fmol/ml in plasma and 5–10 fmol/g in tissue samples)<sup>157–162</sup>. Although RIA and ELISA are the traditional methods for quantifying angiotensin peptides given their high sensitivity and specificity, these methods rely heavily on the characteristics of the antibodies. Therefore, liquid chromatography–mass spectrometry is a promising approach for obtaining reliable read-outs, given its capacity to detect renin–angiotensin system (RAS) peptides by assessing their unique mass-to-charge spectra, which might also allow measurement of potential post-translational modifications in these peptides<sup>163</sup>. However, this technique is not without drawbacks. A complex spectrum resulting from measurement of multiple components with similar mass-to-charge ratios has been reported<sup>164</sup>. Moreover, this approach is expensive and requires very specific expertise<sup>163</sup>. These considerations are of paramount importance, given that inaccurate measurement of RAS peptides can lead to erroneous conclusions that might cloud our understanding of the non-canonical RAS.



Therefore, the identification of these peptides requires the use of high-precision approaches, such as high-performance liquid chromatography and mass spectrometry (BOX 2). Furthermore, one of the fundamental problems associated with the use of these peptides in the clinical context is their short plasma half-life, owing to rapid enzymatic degradation. In each of the numerous ongoing clinical trials assessing the effects of angiotensin 1–7 in CVDs, angiotensin 1–7 is administered via subcutaneous or intravenous injection<sup>126–131</sup>. However, a cyclized angiotensin 1–7 analogue has been described that has increased half-life, improved resistance to enzymatic degradation and superior functional activity compared with natural angiotensin 1–7<sup>136</sup>. Similar chemical modifications to the angiotensin 1–9 peptide might also prolong the half-life of the peptide. However, the non-peptide agonist C21, which has a half-life of 4–6 h, can also induce AT<sub>2</sub>R activation<sup>137</sup>. This agonist has high selectivity for its receptor and is well tolerated<sup>137,138</sup>. However, although the results to date are promising, angiotensin 1–9 still requires extensive safety and efficacy assessment as a potential endogenous AT<sub>2</sub>R agonist.

The oral bioavailability of C21 is only 30%, and this agonist has also been reported to modulate epigenetic mechanisms associated with the pathophysiology of diabetic nephropathy<sup>137</sup>, raising the possibility of unwanted off-target effects if used to treat CVD. Studies comparing the effects of C21 and angiotensin 1–9 will be useful to establish the potential differences between the two agents.

### Complementary agents

Once the challenges hindering clinical translation of counter-regulatory RAS components for the treatment of CVD have been overcome, these therapeutic agents might be used to complement traditional pharmacological treatments. Such complementary drugs are necessary, because even gold-standard drugs for hypertension are associated with issues such as suboptimal drug efficacy and adherence. Most patients with hypertension, especially those with comorbidities, require two or more drugs to manage their blood pressure levels<sup>139,140</sup>. Furthermore, many of these patients require two or more doses each day<sup>140</sup>, suggesting that the separate use of ACE inhibitors or angiotensin II receptor antagonists is not always effective. The use of more than one drug and the need for multiple doses per day can increase the incidence of adverse events, which can result in loss of adherence<sup>140,141</sup>. In addition, a longitudinal study that evaluated the dosing histories of 4,783 patients taking antihypertensive drugs found that nearly half of the patient cohort discontinued the treatment<sup>142</sup>, which results in poorly controlled hypertension<sup>143</sup>. Combining these counter-regulatory RAS peptides with the current gold-standard antihypertensive drugs in one pill might overcome the need for patients with hypertension and other comorbidities to receive more than one drug or multiple dosages of drugs per day. Counter-regulatory RAS peptides, such as angiotensin 1–7, alamandine or angiotensin 1–9, have been found to be effective in reducing blood pressure and attenuating cardiovascular remodelling in preclinical studies<sup>67,73,144</sup>. These effects might be achieved with fewer adverse reactions in patients with hypertension compared with current antihypertensive therapies, which in turn might improve treatment adherence. Combining angiotensin 1–7 with the angiotensin-receptor blocker losartan might increase or extend its blood pressure-lowering capacity<sup>145</sup>. Importantly, the anti-atherosclerotic effects of dual angiotensin 1–7 and losartan therapy are synergistic<sup>146</sup>. Pharmacological synergy between current gold-standard treatment for CVDs and counter-regulatory RAS peptides might decrease the dosages required to achieve efficacy, thereby reducing adverse effects. However, although the endogenous origin of counter-regulatory RAS components suggests a safe pharmacological profile, the current lack of robust evidence in patients means that this hypothesis remains to be tested.

### Conclusions

The evidence supporting the protective role of the counter-regulatory RAS in CVD is robust but incomplete. In addition to the methodological pitfalls that must be overcome, future research should also be

conducted in large animals with high translational value to further confirm the data from the studies carried out in vitro and in small-animal models. The roles of other RAS peptides, such as angiotensin III and angiotensin IV, in the cardiovascular system warrant further investigation. Furthermore, the assessment of classical and counter-regulatory RAS peptides during routine clinical evaluation in patients with CVD should be considered, although development of practical, affordable and accurate methods to assess these levels are required

to achieve reliable readouts. The balance — or imbalance — of the levels of these peptides in plasma or urine might be useful as markers of CVD. Moreover, a thorough evaluation of the counter-regulatory RAS profile of each patient might bring current therapeutic approaches a step closer to the goal of precision medicine, allowing tailored treatment plans for each patient to optimize drug efficacy and adherence.

Published online 19 August 2019

1. Ferrario, C. M. Role of angiotensin II in cardiovascular disease therapeutic implications of more than a century of research. *J. Renin Angiotensin Aldosterone Syst.* **7**, 3–14 (2006).
2. Karnik, S. S. et al. International Union of Basic and Clinical Pharmacology. XCIX. Angiotensin receptors: interpreters of pathophysiological angiotensinergic stimuli [corrected]. *Pharmacol. Rev.* **67**, 754–819 (2015).
3. Forrester, S. J. et al. Angiotensin II signal transduction: an update on mechanisms of physiology and pathophysiology. *Physiol. Rev.* **98**, 1627–1738 (2018).
4. Teixeira, L. B. et al. Ang-(1-7) is an endogenous beta-arrestin-biased agonist of the AT1 receptor with protective action in cardiac hypertrophy. *Sci. Rep.* **7**, 11903 (2017).
5. Jesus, I. C. G. et al. Alamandine acts via MrgD to induce AMPK/NO activation against ANG II hypertrophy in cardiomyocytes. *Am. J. Physiol. Cell. Physiol.* **314**, C702–C711 (2018).
6. Mendoza-Torres, E. et al. Protection of the myocardium against ischemia/reperfusion injury by angiotensin-(1-9) through an AT2R and Akt-dependent mechanism. *Pharmacol. Res.* **135**, 112–121 (2018).
7. Li, T. et al. Critical role of the chymase/angiotensin-(1-12) axis in modulating cardiomyocyte contractility. *Int. J. Cardiol.* **264**, 137–144 (2018).
8. Yu, L., Yuan, K., Phuong, H. T., Park, B. M. & Kim, S. H. Angiotensin-(1-5), an active mediator of renin-angiotensin system, stimulates ANP secretion via Mas receptor. *Peptides* **86**, 33–41 (2016).
9. Chang, L. et al. Bmal1 in perivascular adipose tissue regulates resting-phase blood pressure through transcriptional regulation of angiotensinogen. *Circulation* **138**, 67–79 (2018).
10. Tetzner, A. et al. G-protein-coupled receptor MrgD is a receptor for angiotensin-(1-7) involving adenylyl cyclase, cAMP, and phosphokinase A. *Hypertension* **68**, 185–194 (2016).
11. Bosnyak, S. et al. Relative affinity of angiotensin peptides and novel ligands at AT1 and AT2 receptors. *Clin. Sci.* **121**, 297–303 (2011).
12. Kostenis, E. et al. G-protein-coupled receptor Mas is a physiological antagonist of the angiotensin II type 1 receptor. *Circulation* **111**, 1806–1813 (2005).
13. Gaidarov, I. et al. Angiotensin (1-7) does not interact directly with MAS1, but can potentially antagonize signaling from the AT1 receptor. *Cell. Signal.* **50**, 9–24 (2018).
14. Santos, R. A. et al. Angiotensin-(1-7) is an endogenous ligand for the G protein-coupled receptor Mas. *Proc. Natl Acad. Sci. USA* **100**, 8258–8263 (2003). **This study shows that angiotensin 1–7 binds to the Mas receptor.**
15. Meems, L. M. G. et al. Design, synthesis, and actions of an innovative bispecific designer peptide. *Hypertension* **73**, 900–909 (2019). **This paper describes the synthesis of a peptide that simultaneously activates the Mas receptor and the particulate guanylyl cyclase A receptor, with strong anti-hypertensive effects.**
16. Leonhardt, J. et al. Evidence for heterodimerization and functional interaction of the angiotensin type 2 receptor and the receptor MAS. *Hypertension* **69**, 1128–1135 (2017).
17. Zhang, H. et al. Structural basis for selectivity and diversity in angiotensin II receptors. *Nature* **544**, 327–332 (2017). **This paper reports the crystal structure of human AT<sub>2</sub>R and evidence showing that this receptor does not bind to G proteins or β-arrestins.**
18. Lobo, M. D., Sobotka, P. A. & Pathak, A. Interventional procedures and future drug therapy for hypertension. *Eur. Heart J.* **38**, 1101–1111 (2017).
19. Pan, X. et al. FGF21 Prevents angiotensin II-induced hypertension and vascular dysfunction by activation of ACE2/angiotensin-(1-7) axis in mice. *Cell Metab.* **27**, 1323–1337.e5 (2018).
20. Lawson, C., Vicencio, J. M., Yellon, D. M. & Davidson, S. M. Microvesicles and exosomes: new players in metabolic and cardiovascular disease. *J. Endocrinol.* **228**, R57–R71 (2016).
21. Yellon, D. M. & Davidson, S. M. Exosomes: nanoparticles involved in cardioprotection? *Circ. Res.* **114**, 325–332 (2014).
22. Pironti, G. et al. Circulating exosomes induced by cardiac pressure overload contain functional angiotensin II type 1 receptors. *Circulation* **131**, 2120–2130 (2015).
23. Lyu, L. et al. A critical role of cardiac fibroblast-derived exosomes in activating renin angiotensin system in cardiomyocytes. *J. Mol. Cell. Cardiol.* **89**, 268–279 (2015).
24. Hamming, I. et al. The emerging role of ACE2 in physiology and disease. *J. Pathol.* **212**, 1–11 (2007).
25. Yamazato, Y. et al. Prevention of pulmonary hypertension by angiotensin-converting enzyme 2 gene transfer. *Hypertension* **54**, 365–371 (2009).
26. Ferreira, A. J. et al. Evidence for angiotensin-converting enzyme 2 as a therapeutic target for the prevention of pulmonary hypertension. *Am. J. Respir. Crit. Care Med.* **179**, 1048–1054 (2009).
27. Li, G. et al. Angiotensin-converting enzyme 2 activation ameliorates pulmonary endothelial dysfunction in rats with pulmonary arterial hypertension through mediating phosphorylation of endothelial nitric oxide synthase. *J. Am. Soc. Hypertens.* **11**, 842–852 (2017).
28. Sztuka, K., Orszulak-Michalak, D. & Jasinska-Stroschein, M. Systematic review and meta-analysis of interventions tested in animal models of pulmonary hypertension. *Vasc. Pharmacol.* **110**, 55–63 (2018).
29. Epelman, S. et al. Soluble angiotensin-converting enzyme 2 in human heart failure: relation with myocardial function and clinical outcomes. *J. Card. Fail.* **15**, 565–571 (2009).
30. Shao, Z. et al. Increasing serum soluble angiotensin-converting enzyme 2 activity after intensive medical therapy is associated with better prognosis in acute decompensated heart failure. *J. Card. Fail.* **19**, 605–610 (2013).
31. Johnson, J. A., West, J., Maynard, K. B. & Hemnes, A. R. ACE2 improves right ventricular function in a pressure overload model. *PLOS ONE* **6**, e20828 (2011).
32. Rathinasabapathy, A. et al. rhACE2 therapy modifies bleomycin-induced pulmonary hypertension via rescue of vascular remodeling. *Front. Physiol.* **9**, 271 (2018).
33. Hemnes, A. R. et al. A potential therapeutic role for angiotensin-converting enzyme 2 in human pulmonary arterial hypertension. *Eur. Respir. J.* **51**, 1702638 (2018).
34. Khan, A. et al. A pilot clinical trial of recombinant human angiotensin-converting enzyme 2 in acute respiratory distress syndrome. *Crit. Care* **21**, 234 (2017).
35. Hampf, V. et al. Intrapulmonary activation of the angiotensin-converting enzyme type 2/angiotensin 1-7/G-protein-coupled Mas receptor axis attenuates pulmonary hypertension in Ren-2 transgenic rats exposed to chronic hypoxia. *Physiol. Res.* **64**, 25–38 (2015).
36. Yamada, K., Iyer, S. N., Chappell, M. C., Ganten, D. & Ferrario, C. M. Converting enzyme determines plasma clearance of angiotensin-(1-7). *Hypertension* **32**, 496–502 (1998).
37. Kastin, A. J. & Pan, W. Concepts for biologically active peptides. *Curr. Pharm. Des.* **16**, 3390–3400 (2010).
38. Bennion, D. M. et al. Neuroprotection by post-stroke administration of an oral formulation of angiotensin-(1-7) in ischaemic stroke. *Exp. Physiol.* **103**, 916–923 (2018).
39. Becker, L. K. et al. Eccentric overload muscle damage is attenuated by a novel angiotensin-(1-7) treatment. *Int. J. Sports Med.* **39**, 743–748 (2018).
40. Sabharwal, R. et al. Chronic oral administration of Ang-(1-7) improves skeletal muscle, autonomic and locomotor phenotypes in muscular dystrophy. *Clin. Sci.* **127**, 101–109 (2014).
41. Bertagnolli, M. et al. An orally active angiotensin-(1-7) inclusion compound and exercise training produce similar cardiovascular effects in spontaneously hypertensive rats. *Peptides* **51**, 65–73 (2014).
42. Marques, F. D. et al. Beneficial effects of long-term administration of an oral formulation of angiotensin-(1-7) in infarcted rats. *Int. J. Hypertens.* **2012**, 795452 (2012).
43. Marques, F. D. et al. An oral formulation of angiotensin-(1-7) produces cardioprotective effects in infarcted and isoproterenol-treated rats. *Hypertension* **57**, 477–483 (2011). **This paper describes a new formulation of angiotensin 1–7 that increases the plasma half-life.**
44. Lula, I. et al. Study of angiotensin-(1-7) vasoactive peptide and its beta-cyclodextrin inclusion complexes: complete sequence-specific NMR assignments and structural studies. *Peptides* **28**, 2199–2210 (2007).
45. Breitling, S. et al. Dose-dependent, therapeutic potential of angiotensin-(1-7) for the treatment of pulmonary arterial hypertension. *Pulm. Circ.* **5**, 649–657 (2015).
46. Malek, V., Sharma, N., Sankrityayan, H. & Gaikwad, A. B. Concurrent neprilysin inhibition and renin-angiotensin system modulations prevented diabetic nephropathy. *Life Sci.* **221**, 159–167 (2019).
47. Ocaranza, M. P. & Jalil, J. E. Protective role of the ACE2/Ang-(1-9) axis in cardiovascular remodeling. *Int. J. Hypertens.* **2012**, 594361 (2012).
48. Cha, S. A., Park, B. M. & Kim, S. H. Angiotensin-(1-9) ameliorates pulmonary arterial hypertension via angiotensin type II receptor. *Korean J. Physiol. Pharmacol.* **22**, 447–456 (2018).
49. Bruce, E. et al. Selective activation of angiotensin AT2 receptors attenuates progression of pulmonary hypertension and inhibits cardiopulmonary fibrosis. *Br. J. Pharmacol.* **172**, 2219–2231 (2015).
50. Wagenaar, G. T. et al. Agonists of MAS oncogene and angiotensin II type 2 receptors attenuate cardiopulmonary disease in rats with neonatal hyperoxia-induced lung injury. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **305**, L341–L351 (2013).
51. Hernandez Prada, J. A. et al. Structure-based identification of small-molecule angiotensin-converting enzyme 2 activators as novel antihypertensive agents. *Hypertension* **51**, 1312–1317 (2008).
52. De Maria, M. L. et al. Anti-hypertensive effects of diminazene aceturate: an angiotensin-converting enzyme 2 activator in rats. *Protein Peptide Lett.* **23**, 9–16 (2016).
53. Wiemer, G., Dobrucki, L. W., Louka, F. R., Malinski, T. & Heitsch, H. AVE 0991, a nonpeptide mimic of the effects of angiotensin-(1-7) on the endothelium. *Hypertension* **40**, 847–852 (2002).
54. Savergnini, S. O. et al. Vascular relaxation, antihypertensive effect, and cardioprotection of a novel peptide agonist of the MAS receptor. *Hypertension* **56**, 112–120 (2010).
55. Liu, P. et al. Novel ACE2-Fc chimeric fusion provides long-lasting hypertension control and organ protection in mouse models of systemic renin angiotensin system activation. *Kidney Int.* **94**, 114–125 (2018).
56. Haber, P. K. et al. Angiotensin-converting enzyme 2-independent action of presumed angiotensin-converting enzyme 2 activators: studies in vivo, ex vivo, and in vitro. *Hypertension* **63**, 774–782 (2014).



57. Shenoy, V. et al. Diminazene attenuates pulmonary hypertension and improves angiogenic progenitor cell functions in experimental models. *Am. J. Respir. Crit. Care Med.* **187**, 648–657 (2013).
58. Singh, Y., Singh, K. & Sharma, P. L. Effect of combination of renin inhibitor and Mas-receptor agonist in DOCA-salt-induced hypertension in rats. *Mol. Cell. Biochem.* **373**, 189–194 (2013).
59. Ma, Y. et al. AVE 0991 attenuates cardiac hypertrophy through reducing oxidative stress. *Biochem. Biophys. Res. Commun.* **474**, 621–625 (2016).
60. Keidar, S., Strizevsky, A., Raz, A. & Gamliel-Lazarovich, A. ACE2 activity is increased in monocyte-derived macrophages from prehypertensive subjects. *Nephrol. Dial. Transplant.* **22**, 597–601 (2007).
61. Ortiz-Perez, J. T. et al. Role of circulating angiotensin converting enzyme 2 in left ventricular remodeling following myocardial infarction: a prospective controlled study. *PLOS ONE* **8**, e61695 (2013).
62. Soro-Paavonen, A. et al. Circulating ACE2 activity is increased in patients with type 1 diabetes and vascular complications. *J. Hypertens.* **30**, 375–383 (2012).
63. Roberts, M. A., Velkoska, E., Ierino, F. L. & Burrell, L. M. Angiotensin-converting enzyme 2 activity in patients with chronic kidney disease. *Nephrol. Dial. Transplant.* **28**, 2287–2294 (2013).
64. Khlestova, G. V. et al. Dynamics of renin, angiotensin II, and angiotensin (1-7) during pregnancy and predisposition to hypertension-associated complications. *Bull. Exp. Biol. Med.* **165**, 438–439 (2018).
65. Ferrario, C. M. et al. Characterization of angiotensin-(1-7) in the urine of normal and essential hypertensive subjects. *Am. J. Hypertens.* **11**, 137–146 (1998).
66. Schinzari, F. et al. Favorable vascular actions of angiotensin-(1-7) in human obesity. *Hypertension* **71**, 185–191 (2018).
67. Lautner, R. Q. et al. Discovery and characterization of alamandine: a novel component of the renin-angiotensin system. *Circ. Res.* **112**, 1104–1111 (2013).
68. Soltani Hekmat, A., Javanmardi, K., Kouhpayeh, A., Baharamali, E. & Farjam, M. Differences in cardiovascular responses to alamandine in two-kidney, one clip hypertensive and normotensive rats. *Circ. J.* **81**, 405–412 (2017).
69. Souza-Neto, F. P. et al. Alamandine attenuates arterial remodelling induced by transverse aortic constriction in mice. *Clin. Sci.* **133**, 629–643 (2019).
70. Ichiki, T. et al. Effects on blood pressure and exploratory behaviour of mice lacking angiotensin II type 2 receptor. *Nature* **377**, 748–750 (1995).
71. Hein, L., Barsh, G. S., Pratt, R. E., Dzau, V. J. & Kobilka, B. K. Behavioural and cardiovascular effects of disrupting the angiotensin II type 2 receptor in mice. *Nature* **377**, 744–747 (1995).
72. Tsutsumi, Y. et al. Angiotensin II type 2 receptor overexpression activates the vascular kinin system and causes vasodilation. *J. Clin. Invest.* **104**, 925–935 (1999).
73. Ocaranza, M. P. et al. Angiotensin-(1-9) reverses experimental hypertension and cardiovascular damage by inhibition of the angiotensin converting enzyme/Ang II axis. *J. Hypertens.* **32**, 771–783 (2014).
74. Ali, Q., Wu, Y. & Hussain, T. Chronic AT2 receptor activation increases renal ACE2 activity, attenuates AT1 receptor function and blood pressure in obese Zucker rats. *Kidney Int.* **84**, 931–939 (2013).
75. Li, X. C. & Widdop, R. E. AT2 receptor-mediated vasodilatation is unmasked by AT1 receptor blockade in conscious SHR. *Br. J. Pharmacol.* **142**, 821–830 (2004).
76. Benndorf, R., Boger, R. H., Ergun, S., Steenpass, A. & Wieland, T. Angiotensin II type 2 receptor inhibits vascular endothelial growth factor-induced migration and in vitro tube formation of human endothelial cells. *Circ. Res.* **93**, 438–447 (2003).
77. Ocaranza, M. P. et al. Rho kinase inhibition activates the homologous angiotensin-converting enzyme-angiotensin-(1-9) axis in experimental hypertension. *J. Hypertens.* **29**, 706–715 (2011).
78. Xu, X. et al. RhoA-Rho associated kinase signaling leads to renin-angiotensin system imbalance and angiotensin converting enzyme 2 has a protective role in acute pulmonary embolism. *Thromb. Res.* **176**, 85–94 (2019).
79. Fattah, C. et al. Gene therapy with angiotensin-(1-9) preserves left ventricular systolic function after myocardial infarction. *J. Am. Coll. Cardiol.* **68**, 2652–2666 (2016).
- This study showed that the use of gene therapy to overexpress angiotensin 1–9 prevents cardiac dysfunction and improves survival after myocardial infarction in mice.**
80. Flores-Munoz, M. et al. Angiotensin-(1-9) attenuates cardiac fibrosis in the stroke-prone spontaneously hypertensive rat via the angiotensin type 2 receptor. *Hypertension* **59**, 300–307 (2012).
- Angiotensin 1–9 reduces hypertensive cardiovascular remodelling.**
81. Reddy, R. et al. Circulating angiotensin peptides levels in acute respiratory distress syndrome correlate with clinical outcomes: a pilot study. *PLOS ONE* **14**, e0213096 (2019).
82. Crackower, M. A. et al. Angiotensin-converting enzyme 2 is an essential regulator of heart function. *Nature* **417**, 822–828 (2002).
- This paper is the first publication on the role of ACE2 in heart function.**
83. Rentsch, B. et al. Transgenic angiotensin-converting enzyme 2 overexpression in vessels of SHRSP rats reduces blood pressure and improves endothelial function. *Hypertension* **52**, 967–973 (2008).
84. Xu, P., Sriramula, S. & Lazartigues, E. ACE2/ANG-(1-7)/Mas pathway in the brain: the axis of good. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **300**, R804–R817 (2011).
85. Oudit, G. Y. et al. Angiotensin II-mediated oxidative stress and inflammation mediate the age-dependent cardiomyopathy in ACE2 null mice. *Cardiovasc. Res.* **75**, 29–39 (2007).
86. Epelman, S. et al. Detection of soluble angiotensin-converting enzyme 2 in heart failure: insights into the endogenous counter-regulatory pathway of the renin-angiotensin-aldosterone system. *J. Am. Coll. Cardiol.* **52**, 750–754 (2008).
87. Redfield, M. M. Heart failure with preserved ejection fraction. *N. Engl. J. Med.* **375**, 1868–1877 (2016).
88. Zhong, J. et al. Angiotensin-converting enzyme 2 suppresses pathological hypertrophy, myocardial fibrosis, and cardiac dysfunction. *Circulation* **122**, 717–728 (2010).
89. Oliveira, A. C. et al. Genetic deletion of the alamandine receptor MRGD leads to dilated cardiomyopathy in mice. *Am. J. Physiol. Heart Circ. Physiol.* **316**, H123–H135 (2019).
90. Gao, J., Zucker, I. H. & Gao, L. Activation of central angiotensin type 2 receptors by compound 21 improves arterial baroreflex sensitivity in rats with heart failure. *Am. J. Hypertens.* **27**, 1248–1256 (2014).
91. Burrell, L. M. et al. Myocardial infarction increases ACE2 expression in rat and humans. *Eur. Heart J.* **26**, 369–375; discussion 322–364 (2005).
92. Kassiri, Z. et al. Loss of angiotensin-converting enzyme 2 accelerates maladaptive left ventricular remodeling in response to myocardial infarction. *Circ. Heart Fail.* **2**, 446–455 (2009).
93. Der Sarkissian, S. et al. Cardiac overexpression of angiotensin converting enzyme 2 protects the heart from ischemia-induced pathophysiology. *Hypertension* **51**, 712–718 (2008).
94. Santos, R. A. et al. Expression of an angiotensin-(1-7)-producing fusion protein produces cardioprotective effects in rats. *Physiol. Genomics* **17**, 292–299 (2004).
95. Oishi, Y. et al. Cardioprotective role of AT2 receptor in postinfarction left ventricular remodeling. *Hypertension* **41**, 814–818 (2003).
96. Brede, M. et al. Cardiac hypertrophy is associated with decreased eNOS expression in angiotensin AT2 receptor-deficient mice. *Hypertension* **42**, 1177–1182 (2003).
97. Adachi, Y. et al. Angiotensin II type 2 receptor deficiency exacerbates heart failure and reduces survival after acute myocardial infarction in mice. *Circulation* **107**, 2406–2408 (2003).
98. Yang, Z. et al. Angiotensin II type 2 receptor overexpression preserves left ventricular function after myocardial infarction. *Circulation* **106**, 106–111 (2002).
99. Qi, Y. et al. Moderate cardiac-selective overexpression of angiotensin II type 2 receptor protects cardiac functions from ischaemic injury. *Exp. Physiol.* **97**, 89–101 (2012).
100. Lauer, D. et al. Angiotensin type 2 receptor stimulation ameliorates left ventricular fibrosis and dysfunction via regulation of tissue inhibitor of matrix metalloproteinase 1/matrix metalloproteinase 9 axis and transforming growth factor beta1 in the rat heart. *Hypertension* **63**, e60–e67 (2014).
101. Ruparel, N., Chai, J. T., Fisher, E. A. & Choudhury, R. P. Inflammatory processes in cardiovascular disease: a route to targeted therapies. *Nat. Rev. Cardiol.* **14**, 133–144 (2017).
102. Shirazi, L. F., Bissett, J., Romeo, F. & Mehta, J. L. Role of inflammation in heart failure. *Curr. Atheroscler. Rep.* **19**, 27 (2017).
103. Golia, E. et al. Inflammation and cardiovascular disease: from pathogenesis to therapeutic target. *Curr. Atheroscler. Rep.* **16**, 435 (2014).
104. Guzik, T. J. & Touyz, R. M. Oxidative stress, inflammation, and vascular aging in hypertension. *Hypertension* **70**, 660–667 (2017).
105. McMaster, W. G., Kirabo, A., Madhur, M. S. & Harrison, D. G. Inflammation, immunity, and hypertensive end-organ damage. *Circ. Res.* **116**, 1022–1033 (2015).
106. Hoch, N. E. et al. Regulation of T-cell function by endogenously produced angiotensin II. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **296**, R208–R216 (2009).
107. Jurewicz, M. et al. Human T and natural killer cells possess a functional renin-angiotensin system: further mechanisms of angiotensin II-induced inflammation. *J. Am. Soc. Nephrol.* **18**, 1093–1102 (2007).
108. Dinh, Q. N., Drummond, G. R., Sobey, C. G. & Chrissobolis, S. Roles of inflammation, oxidative stress, and vascular dysfunction in hypertension. *Biomed. Res. Int.* **2014**, 406960 (2014).
109. Rutkowska-Zapala, M. et al. Human monocyte subsets exhibit divergent angiotensin I-converting activity. *Clin. Exp. Immunol.* **181**, 126–132 (2015).
110. da Silveira, K. D. et al. Anti-inflammatory effects of the activation of the angiotensin-(1-7) receptor, MAS, in experimental models of arthritis. *J. Immunol.* **185**, 5569–5576 (2010).
111. Oliveira-Lima, O. C. et al. Mas receptor deficiency exacerbates lipopolysaccharide-induced cerebral and systemic inflammation in mice. *Immunobiology* **220**, 1311–1321 (2015).
112. Passos-Silva, D. G., Verano-Braga, T. & Santos, R. A. Angiotensin-(1-7): beyond the cardio-renal actions. *Clin. Sci.* **124**, 443–456 (2013).
113. Magalhaes, G. S. et al. Angiotensin-(1-7) attenuates airway remodelling and hyperresponsiveness in a model of chronic allergic lung inflammation. *Br. J. Pharmacol.* **172**, 2330–2342 (2015).
114. Jawien, J. et al. Angiotensin-(1-7) receptor Mas agonist ameliorates progress of atherosclerosis in apoE-knockout mice. *J. Physiol. Pharmacol.* **63**, 77–85 (2012).
115. Tesanovic, S., Vinh, A., Gaspari, T. A., Casley, D. & Widdop, R. E. Vasoprotective and atheroprotective effects of angiotensin (1-7) in apolipoprotein E-deficient mice. *Arterioscler. Thromb. Vasc. Biol.* **30**, 1606–1613 (2010).
116. Magalhaes, G. S. et al. Angiotensin-(1-7) promotes resolution of eosinophilic inflammation in an experimental model of asthma. *Front. Immunol.* **9**, 58 (2018).
117. Dhande, I., Ma, W. & Hussain, T. Angiotensin AT2 receptor stimulation is anti-inflammatory in lipopolysaccharide-activated THP-1 macrophages via increased interleukin-10 production. *Hypertens. Res.* **38**, 21–29 (2015).
118. Dhande, I., Ali, Q. & Hussain, T. Proximal tubule angiotensin AT2 receptors mediate an anti-inflammatory response via interleukin-10: role in renoprotection in obese rats. *Hypertension* **61**, 1218–1226 (2013).
119. Kaschina, E. et al. Angiotensin II type 2 receptor stimulation: a novel option of therapeutic interference with the renin-angiotensin system in myocardial infarction? *Circulation* **118**, 2523–2532 (2008).
120. Gonzalez, L. et al. Angiotensin-(1-9) reduces cardiovascular and renal inflammation in experimental renin-independent hypertension. *Biochem. Pharmacol.* **156**, 357–370 (2018).
- This paper describes that angiotensin 1–9 is an anti-inflammatory agent.**
121. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT00886553> (2009).
122. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT00771810> (2017).
123. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT01884051> (2019).
124. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT03177603> (2019).

125. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT03000686> (2019).
126. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT03252093> (2019).
127. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT02245230> (2018).
128. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT02591173> (2018).
129. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT02646475> (2018).
130. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT03159988> (2019).
131. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT03615196> (2019).
132. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT03240068> (2019).
133. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT03604289> (2019).
134. Basu, R. et al. Roles of angiotensin peptides and recombinant human ACE2 in heart failure. *J. Am. Coll. Cardiol.* **69**, 805–819 (2017).
135. Ocaranza, M. P. et al. Angiotensin-(1-9) regulates cardiac hypertrophy in vivo and in vitro. *J. Hypertens.* **28**, 1054–1064 (2010). **This paper describes for the first time the anti-hypertrophic effect of angiotensin 1–9.**
136. Kluskens, L. D. et al. Angiotensin-(1-7) with thioether bridge: an angiotensin-converting enzyme-resistant, potent angiotensin-(1-7) analog. *J. Pharmacol. Exp. Ther.* **328**, 849–854 (2009).
137. Pandey, A. & Gaikwad, A. B. AT2 receptor agonist compound 21: a silver lining for diabetic nephropathy. *Eur. J. Pharmacol.* **815**, 251–257 (2017).
138. Wan, Y. et al. Design, synthesis, and biological evaluation of the first selective nonpeptide AT2 receptor agonist. *J. Med. Chem.* **47**, 5995–6008 (2004).
139. Kalra, S., Kalra, B. & Agrawal, N. Combination therapy in hypertension: an update. *Diabetol. Metab. Syndr.* **2**, 44 (2010).
140. Gradman, A. H., Basile, J. N., Carter, B. L., Bakris, G. L. & American Society of Hypertension Writing Group. Combination therapy in hypertension. *J. Am. Soc. Hypertension* **4**, 42–50 (2010).
141. Gebreyohannes, E. A., Bhagavathula, A. S., Abebe, T. B., Tefera, Y. G. & Abegaz, T. M. Adverse effects and non-adherence to antihypertensive medications in University of Gondar Comprehensive Specialized Hospital. *Clin. Hypertens.* **25**, 1 (2019).
142. Vrijens, B., Vincze, G., Kristanto, P., Urquhart, J. & Burnier, M. Adherence to prescribed antihypertensive drug treatments: longitudinal study of electronically compiled dosing histories. *BMJ* **336**, 1114–1117 (2008).
143. Burnier, M. & Egan, B. M. Adherence in hypertension. *Circ. Res.* **124**, 1124–1140 (2019).
144. Grobe, J. L. et al. Prevention of angiotensin II-induced cardiac remodeling by angiotensin-(1-7). *Am. J. Physiol. Heart Circ. Physiol.* **292**, H736–H742 (2007).
145. Machado-Silva, A., Passos-Silva, D., Santos, R. A. & Sinisterra, R. D. Therapeutic uses for angiotensin-(1-7). *Expert Opin. Ther. Pat.* **26**, 669–678 (2016).
146. Yang, J. et al. Comparison of angiotensin-(1-7), losartan and their combination on atherosclerotic plaque formation in apolipoprotein E knockout mice. *Atherosclerosis* **240**, 544–549 (2015).
147. Young, D., Waitches, G., Birchmeier, C., Fasano, O. & Wigler, M. Isolation and characterization of a new cellular oncogene encoding a protein with multiple potential transmembrane domains. *Cell* **45**, 711–719 (1986).
148. Johnson, H. & Drummer, O. H. Hydrolysis of angiotensin I by peptidases in homogenates of rat lung and aorta. *Biochem. Pharmacol.* **37**, 1131–1136 (1988).
149. Santos, R. A. et al. Converting enzyme activity and angiotensin metabolism in the dog brainstem. *Hypertension* **11**, 1153–1157 (1988).
150. Campagnole-Santos, M. J. et al. Cardiovascular effects of angiotensin-(1-7) injected into the dorsal medulla of rats. *Am. J. Physiol.* **257**, H324–H329 (1989).
151. Tipnis, S. R. et al. A human homolog of angiotensin-converting enzyme. Cloning and functional expression as a captopril-insensitive carboxypeptidase. *J. Biol. Chem.* **275**, 33238–33243 (2000).
152. Donoghue, M. et al. A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9. *Circulation Res.* **87**, E1–E9 (2000).
153. Ferreira, A. J., Santos, R. A. & Almeida, A. P. Angiotensin-(1-7): cardioprotective effect in myocardial ischemia/reperfusion. *Hypertension* **38**, 665–668 (2001).
154. US National Library of Medicine. *ClinicalTrials.gov* <https://clinicaltrials.gov/ct2/show/NCT00471562> (2017).
155. Liu, C. et al. Alamandine attenuates hypertension and cardiac hypertrophy in hypertensive rats. *Amino Acids* **50**, 1071–1081 (2018).
156. Park, B. M., Phuong, H. T. A., Yu, L. & Kim, S. H. Alamandine protects the heart against reperfusion injury via the MrgD receptor. *Circ. J.* **82**, 2584–2593 (2018).
157. Danser, A. H. et al. Cardiac renin and angiotensins. Uptake from plasma versus in situ synthesis. *Hypertension* **24**, 37–48 (1994).
158. Lawrence, A. C., Evin, G., Kladis, A. & Campbell, D. J. An alternative strategy for the radioimmunoassay of angiotensin peptides using amino-terminal-directed antisera: measurement of eight angiotensin peptides in human plasma. *J. Hypertens.* **8**, 715–724 (1990).
159. Alexiou, T. et al. Angiotensinogen and angiotensin-converting enzyme gene copy number and angiotensin and bradykinin peptide levels in mice. *J. Hypertens.* **23**, 945–954 (2005).
160. Ocaranza, M. P. et al. Enalapril attenuates downregulation of angiotensin-converting enzyme 2 in the late phase of ventricular dysfunction in myocardial infarcted rat. *Hypertension* **48**, 572–578 (2006). **This study is the first to show that angiotensin 1–9 counteracts the pathophysiological effects of angiotensin II.**
161. Sharp, S., Poglitsch, M., Zilla, P., Davies, N. H. & Sturrock, E. D. Pharmacodynamic effects of C-domain-specific ACE inhibitors on the renin-angiotensin system in myocardial infarcted rats. *J. Renin Angiotensin Aldosterone Syst.* **16**, 1149–1158 (2015).
162. Haschke, M. et al. Pharmacokinetics and pharmacodynamics of recombinant human angiotensin-converting enzyme 2 in healthy human subjects. *Clin. Pharmacokinet.* **52**, 783–792 (2013).
163. Chappell, M. C. Biochemical evaluation of the renin-angiotensin system: the good, bad, and absolute? *Am. J. Physiol. Heart Circ. Physiol.* **310**, H137–H152 (2016).
164. Lortie, M., Bark, S., Blantz, R. & Hook, V. Detecting low-abundance vasoactive peptides in plasma: progress toward absolute quantitation using nano liquid chromatography-mass spectrometry. *Anal. Biochem.* **394**, 164–170 (2009).
165. Aguilera, G. Role of angiotensin II receptor subtypes on the regulation of aldosterone secretion in the adrenal glomerulosa zone in the rat. *Mol. Cell. Endocrinol.* **90**, 53–60 (1992).
166. Qadri, F. et al. Angiotensin II-induced vasopressin release is mediated through alpha-1 adrenoceptors and angiotensin II AT1 receptors in the supraoptic nucleus. *J. Pharmacol. Exp. Ther.* **267**, 567–574 (1993).
167. Huang, B. S., Chen, A., Ahmad, M., Wang, H. W. & Leenen, F. H. Mineralocorticoid and AT1 receptors in the paraventricular nucleus contribute to sympathetic hyperactivity and cardiac dysfunction in rats post myocardial infarct. *J. Physiol.* **592**, 3273–3286 (2014).
168. Iyer, S. N., Lu, D., Katovich, M. J. & Raizada, M. K. Chronic control of high blood pressure in the spontaneously hypertensive rat by delivery of angiotensin type 1 receptor antisense. *Proc. Natl Acad. Sci. USA* **93**, 9960–9965 (1996).
169. Li, Q., Feenstra, M., Pfaffendorf, M., Eijssman, L. & van Zwieten, P. A. Comparative vasoconstrictor effects of angiotensin II, III, and IV in human isolated saphenous vein. *J. Cardiovasc. Pharmacol.* **29**, 451–456 (1997).
170. Sadoshima, J. & Izumo, S. Molecular characterization of angiotensin II-induced hypertrophy of cardiac myocytes and hyperplasia of cardiac fibroblasts. Critical role of the AT1 receptor subtype. *Circ. Res.* **73**, 413–423 (1993).
171. Schieffer, B. et al. Comparative effects of chronic angiotensin-converting enzyme inhibition and angiotensin II type 1 receptor blockade on cardiac remodeling after myocardial infarction in the rat. *Circulation* **89**, 2273–2282 (1994).
172. Wolf, G. et al. Angiotensin II activates nuclear transcription factor-kappaB through AT1 and AT2 receptors. *Kidney Int.* **61**, 1986–1995 (2002).
173. Viswanathan, M., Stromberg, C., Seltzer, A. & Saavedra, J. M. Balloon angioplasty enhances the expression of angiotensin II AT1 receptors in neointima of rat aorta. *J. Clin. Invest.* **90**, 1707–1712 (1992).
174. Jara, Z. P. et al. Tonin overexpression in mice diminishes sympathetic autonomic modulation and alters angiotensin type 1 receptor response. *Front. Med.* **5**, 365 (2018).
175. de Queiroz, T. M., Monteiro, M. M. & Braga, V. A. Angiotensin-II-derived reactive oxygen species on baroreflex sensitivity during hypertension: new perspectives. *Front. Physiol.* **4**, 105 (2013).
176. Kihara, M. et al. Angiotensin II inhibits interleukin-1 beta-induced nitric oxide production in cultured rat mesangial cells. *Kidney Int.* **55**, 1277–1283 (1999).
177. van der Mark, J. & Kline, R. L. Altered pressure natriuresis in chronic angiotensin II hypertension in rats. *Am. J. Physiol.* **266**, R739–R748 (1994).
178. Park, B. M., Cha, S. A., Lee, S. H. & Kim, S. H. Angiotensin IV protects cardiac reperfusion injury by inhibiting apoptosis and inflammation via AT4R in rats. *Peptides* **79**, 66–74 (2016).
179. Handa, R. K., Krebs, L. T., Harding, J. W. & Handa, S. E. Angiotensin IV AT4-receptor system in the rat kidney. *Am. J. Physiol.* **274**, F290–F299 (1998).
180. Kramar, E. A., Krishnan, R., Harding, J. W. & Wright, J. W. Role of nitric oxide in angiotensin IV-induced increases in cerebral blood flow. *Regul. Pept.* **74**, 185–192 (1998).
181. Coleman, J. K. et al. Autoradiographic identification of kidney angiotensin IV binding sites and angiotensin IV-induced renal cortical blood flow changes in rats. *Peptides* **19**, 269–277 (1998).
182. Qiu, H. et al. Effect of berberine on PPARalpha-NO signalling pathway in vascular smooth muscle cell proliferation induced by angiotensin IV. *Pharm. Biol.* **55**, 227–232 (2017).
183. Padia, S. H. et al. Conversion of renal angiotensin II to angiotensin III is critical for AT2 receptor-mediated natriuresis in rats. *Hypertension* **51**, 460–465 (2008).
184. Fontes, M. A. et al. Evidence that angiotensin-(1-7) plays a role in the central control of blood pressure at the ventro-lateral medulla acting through specific receptors. *Brain Res.* **665**, 175–180 (1994).
185. Xia, H. & Lazartigues, E. Angiotensin-converting enzyme 2: central regulator for cardiovascular function. *Curr. Hypertens. Rep.* **12**, 170–175 (2010).
186. Li, P., Chappell, M. C., Ferrario, C. M. & Brosnihan, K. B. Angiotensin-(1-7) augments bradykinin-induced vasodilation by competing with ACE and releasing nitric oxide. *Hypertension* **29**, 394–400 (1997).
187. DelliPizzi, A. M., Hilchey, S. D. & Bell-Quilley, C. P. Natriuretic action of angiotensin(1-7). *Br. J. Pharmacol.* **111**, 1–3 (1994).
188. Garcia-Espinosa, M. A., Shaltout, H. A., Gallagher, P. E., Chappell, M. C. & Diz, D. I. In vivo expression of angiotensin-(1-7) lowers blood pressure and improves baroreflex function in transgenic (mRen2)27 rats. *J. Cardiovasc. Pharmacol.* **60**, 150–157 (2012).
189. Sakima, A. et al. Baroreceptor reflex regulation in anesthetized transgenic rats with low glia-derived angiotensinogen. *Am. J. Physiol. Heart Circ. Physiol.* **292**, H1412–H1419 (2007).
190. Soares, E. R., Barbosa, C. M., Campagnole-Santos, M. J., Santos, R. A. S. & Alzamora, A. C. Hypotensive effect induced by microinjection of alamandine, a derivative of angiotensin-(1-7), into caudal ventrolateral medulla of 2K1C hypertensive rats. *Peptides* **96**, 67–75 (2017).
191. AbdAlla, S., Lother, H., Abdel-tawab, A. M. & Quitterer, U. The angiotensin II AT2 receptor is an AT1 receptor antagonist. *J. Biol. Chem.* **276**, 39721–39726 (2001).
192. Bedecs, K. et al. Angiotensin II type 2 receptors mediate inhibition of mitogen-activated protein kinase cascade and functional activation of SHP-1 tyrosine phosphatase. *Biochem. J.* **325**, 449–454 (1997).
193. Horiuchi, M., Akishima, M. & Dzau, V. J. Molecular and cellular mechanism of angiotensin II-mediated apoptosis. *Endocr. Res.* **24**, 307–314 (1998).
194. Senbonmatsu, T. et al. A novel angiotensin II type 2 receptor signaling pathway: possible role in cardiac hypertrophy. *EMBO J.* **22**, 6471–6482 (2003).
195. Gao, S. et al. Angiotensin AT2 receptor agonist stimulates high stretch induced- ANP secretion via PI3K/NO/sGC/PKG/pathway. *Peptides* **47**, 36–44 (2013).



196. Cha, S. A., Park, B. M., Gao, S. & Kim, S. H. Stimulation of ANP by angiotensin-(1-9) via the angiotensin type 2 receptor. *Life Sci.* **93**, 934–940 (2013).
197. Abadir, P. M., Periasamy, A., Carey, R. M. & Siragy, H. M. Angiotensin II type 2 receptor-bradykinin B2 receptor functional heterodimerization. *Hypertension* **48**, 316–322 (2006).
198. Tao, X. et al. Angiotensin-(1-7) attenuates angiotensin II-induced signalling associated with activation of a tyrosine phosphatase in Sprague-Dawley rats cardiac fibroblasts. *Biol. Cell.* **106**, 182–192 (2014).
199. McCollum, L. T., Gallagher, P. E. & Tallant, E. A. Angiotensin-(1-7) abrogates mitogen-stimulated proliferation of cardiac fibroblasts. *Peptides* **34**, 380–388 (2012).
200. Shah, A. et al. Angiotensin-(1-7) stimulates high atrial pacing-induced ANP secretion via Mas/PI3-kinase/Akt axis and Na<sup>+</sup>/H<sup>+</sup> exchanger. *Am. J. Physiol. Heart Circ. Physiol.* **298**, H1365–H1374 (2010).
201. Wang, L. P. et al. Protective role of ACE2-Ang-(1-7)-Mas in myocardial fibrosis by downregulating KCa3.1 channel via ERK1/2 pathway. *Pflug. Arch.* **468**, 2041–2051 (2016).
202. Carver, K. A., Smith, T. L., Gallagher, P. E. & Tallant, E. A. Angiotensin-(1-7) prevents angiotensin II-induced fibrosis in cremaster microvessels. *Microcirculation* **22**, 19–27 (2015).
203. Gomes, E. R. et al. Angiotensin-(1-7) prevents cardiomyocyte pathological remodeling through a nitric oxide/guanosine 3',5'-cyclic monophosphate-dependent pathway. *Hypertension* **55**, 153–160 (2010).

#### Acknowledgements

The authors received funding from Comisión Nacional de Ciencia y Tecnología (CONICYT), Chile (FONDAP 15130011 to M.P.O., J.A.R., L.G., M.C. and S.L.; FONDECYT 1161739 to J.E.J.; FONDECYT 11181000 to J.A.R.; FONDECYT 1140713 to L.G.), Grant Puente Pontificia Universidad

Catolica de Chile (P1705/2017 to M.P.O.), Bayer AG (Program Grants4Targets ID 2017-08-2260 to M.P.O., J.E.J., M.C. and S.L.), and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazil (310515/2015–7 to R.A.S.S.).

#### Author contributions

All authors researched data for the article, contributed to discussion of the content and wrote, reviewed and edited the manuscript before submission.

#### Competing interests

M.P.O., M.C., J.E.J. and S.L. have patents related to the pharmacological effects of angiotensin 1–9. R.A.S.S. has patents related to the pharmacological effects of angiotensin 1–7 and alamandine. J.A.R. and L.G. declare no competing interests.

#### Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.