Mohamed Khider University of Biskra Faculty of Exact Sciences and Natural and Life Sciences

1st year LMD – SNV Biology Subject: Chemistry 2 Academic year: 2023/2024

Applied exercises series No. 3

(Solubility - Solubility product)

Exercise 1:

Magnesium hydroxide Mg(OH)2 is a poorly soluble compound, with a molar mass of

58.3 g/mol and which has a solubility product $Ks = 1.2 \ 10^{-11} \ mol^3/mol^3$ at 18 °C.

1. Calculate the solubility of Mg(OH)₂ in grams per liter.

2. Calculate the solubility product of $Ca_3(PO_4)_2$, $S = 2.50 \cdot 10^{-3} \text{ g/l}$; M = 310 g/mol.

Exercise 2:

Bladder (urinary) stones are made up of calcium oxalate, CaC₂O₄. Calcium oxalate is an ionic solid that is poorly soluble in water.

1. Calculate the molar solubility and mass solubility of CaC₂O₄.

2. What minimum volume of pure water should be used to dissolve a 1.1 g urinary stone?

3. The water used actually contains calcium chloride at a concentration of 10^{-4} mol/l. In this case, what is the minimum volume of water necessary to dissolve the same urinary stone of 1.1 g?

<u>We give</u>: the solubility product of calcium oxalate: $Ks = 10^{-8.6}$.

Exercise 3:

The solubility product of lead sulfate $PbSO_4$ is $10^{-8.6}$. Calculate its molar and mass solubility in:

1. Pure water.

2. A solution of lead nitrate $Pb(NO_3)_2$ 0.10 mol/l. calculate the new solubility S', what can we conclude?

Mohamed Khider University of Biskra Faculty of Exact Sciences and Natural and Life Sciences

1st year LMD – SNV Biology Subject: Chemistry 2 Academic year: 2023/2024

Applied exercises series No. 4

(Oxidation-reduction reaction)

Exercise 1:

• Calculate the degrees (numbers) of oxidation of the underlined atoms:

K<u>Mn</u>O₄, H<u>N</u>O₃, H₃PO₄, Cr₂O₇⁻², Na<u>H</u>, O₂, F₂O, I⁻.

Exercise 2:

1. Equilibrium the following reaction in an acidic medium:

 $Cr_2O_7^{-2} + H_2SO_3 \rightarrow Cr^{+3} + HSO_4^{-1}$

2. Equilibrium the following reaction in an acidic medium:

 $NO_3^- + HS^- \longrightarrow NO_2^- + S$

Exercise 3:

1. Express the electrode potential of the Riboflavin-Leucoriboflavin (Rb/RbH_2) system as a function of the pH of the solution at 25 °C.

2. Calculate the equilibrium constant K of the torque equation:

 $(Fe^{3+}/Fe^{2+}) \rightarrow E10 = 0.771 \text{ V}.$

 $(\text{Sn}^{4+}/\text{Sn}^{2+}) \rightarrow \text{E20} = 0.14 \text{ V}.$

Exercise 4:

• What is the e.m.f. from the following battery?

Ag/Ag⁺ (10⁻¹ M) // Cu²⁺ (10⁻² M)/Cu

We give: the standard potentials of the two couples:

 $E^{0}(Ag^{+}/Ag) = 0.80 V, \quad E^{0}(Cu^{2+}/Cu) = 0.34 V$